iv) naphthyl,

5

15

20

25

- v) furanyl, oxazolyl, isoxazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, thiophenyl, thiazolyl, isothiazolyl, pyrrolyl, imidazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 3-indoxazinyl, 2-benzoxazolyl, 2- or 3-benzothiophenyl, 2- or 3-benzofuranyl, 2- or 3-indolyl, 2-benzthiazolyl, 2-benzimidazolyl, 3-indazolyl, and
- vi) pyridinyl, pyridinyl-N-oxide, pyrazinyl, pyrimidinyl, pyridazinyl, 1-, 3- or 4- isoquinolinyl, 2-, 3- or 4-quinolinyl, 2- or 3-quinoxalinyl, 2- or 4-quinazolinyl,
- 10 8. The compound of claim 1 wherein R<sup>2</sup>, optionally substituted with R<sup>q</sup>, is selected from the group consisting of phenyl, naphthalenyl, pyridinyl, thiophenyl, benzothiophenyl, furanyl, benzofuranyl, indolyl, indolinyl, isoquinolinyl and quinolinyl.

The compound of claim 1 wherein R<sup>2</sup> is selected from the group consisting of 4-methyl-phenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 3,4-dichloro-phenyl, benzo[1,3]dioxol-5-yl, 2,3-diydro benzo[1,4]dioxin-6-yl, 4-methoxy-phenyl, phenyl, 4-phenoxy-phenyl, naphthalen-2-yl, pyridin-3-yl, 2-chloro-pyridin-3-yl, pyridin-4-ylmethyl, 4-benzyloxy-phenyl, 4-dimethylamino-phenyl, 4-bromo-3-methyl-phenyl, 3-methoxy-4-methyl-phenyl,

- 3-cyclopentyloxy-4-methoxy-phenyl, 4-bromo-2-chloro-phenyl, 4-bromo-phenyl, 3-dimethylamino-phenyl, 4-morpholin-1-yl-phenyl, 4-pyrrolidin-1-yl-phenyl, 4-(N-propylamino)-phenyl, 4-(N-isobutylamino)-phenyl, 4-diethylamino-phenyl, 4-(N-allylamino)-phenyl, 4-(N-isopropylamino)-phenyl, 4-(N-methyl-N-propylamino)-phenyl, 4-(N-methyl-N-isopropylamino)-phenyl, 4-(N-methyl-N-isopr
- ethylamino)-phenyl, 4-(N-methyl-N-propylamino)-phenyl, 4-(N-methyl-N-propylamino)-2-chloro-phenyl, 4-(N-ethyl-N-methylamino)-2-chloro-phenyl, 4-(pyrrolidin-1-yl)-2-chloro-phenyl, 4-azetidinyl-phenyl, 4-(pyrrolidin-2-one-1-yl)-phenyl, 4-bromo-3-methyl-phenyl, 4-chloro-3-methyl-phenyl, 1-methyl-5-indolinyl, 5-indolinyl, 5-isoquinolinyl, 6-quinolinyl, benzo[1,3]diox-5-yl and 7-methoxy-benzofuran-2-yl.
- The compound of claim 1 wherein R<sup>q</sup> is selected from the group

consisting of -OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, i-propyl, t-butyl, -OCH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -OCH(CH<sub>3</sub>)<sub>2</sub>, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, -Ocyclopentyl,

30